

TECHNICAL RECOMMENDATION

Technical Recommendation

Soybean Spider Mite control with Agri-Mek SC in East Heartland



Source: Elijah Meck



Photo; Syngenta

Application Directions:

- Apply Agri-Mek SC at 1.75-3.5 fl. oz. per acre. Use 1.75 -2.5 fl. oz. per acre for low to moderate infestations and 3.5 fl. oz. per acre for high infestations.
- Apply in a minimum of 10 gallons of water per acre with ground application equipment. Do not apply in less than 5 gallons of water per acre with aircraft. Good coverage of soybean foliage is essential for control of Spider Mites.

Adjuvant Requirement:

- To avoid illegal residues, Agri-Mek SC **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Refer to the **Use Information** section found at the beginning of the Agri-Mek SC label.

Precautions:

- **PHI:** Do not apply within **7 days** of harvest for forage or hay or within **28 days** of harvest for soybean seed. **Do not** allow livestock to graze in treated areas. **Do not** feed treated soybean fodder or silage to meat or dairy animals.
- Do not apply more than 3.5 fl. oz. per acre of Agri-Mek SC or any other foliarly applied abamectin-containing product per application.
- Do not apply more than 7.0 fl. oz. per acre of Agri-Mek SC or any other foliarly applied abamectin-containing product per year. Applications should be at least 7 days apart.
- Do not apply Agri-Mek SC when weather conditions favor drift to aquatic or sensitive areas. Refer to spray drift section on the Agri-Mek SC label.

PJK 06/16

syngenta[®]